

ODD HARMONIOUS LABELING ON SOME STRING GRAPH CLASSES

By Fery Firmansah

Pelabelan Harmonis Ganjil Pada Beberapa Kelas Graf Tali

Odd Harmonious Labeling on Some String Graph Classes

Fery Firmansah*

Prodi Pendidikan Matematika FKIP Universitas Widya Dharma Klaten
Jl. Ki Hajar Dewantara Klaten 57438, Indonesia

e-mail: *feryfirmansah@unwidha.ac.id

Corresponding author*

Abstrak

Graf dengan sifat pelabelan harmonis ganjil disebut sebagai graf harmonis ganjil. Tujuan penelitian ini adalah menemukan sifat pelabelan harmonis ganjil pada beberapa kelas graf tali. Penelitian ini menggunakan metode penelitian kualitatif. Hasil penelitian diperoleh definisi dan konstruksi dari graf tali, gabungan graf tali dan graf tali berlipat. Selanjutnya dibuktikan bahwa graf tali, gabungan graf tali dan graf tali berlipat adalah graf harmonis ganjil.

Kata Kunci : *graf berlipat, graf harmonis ganjil, graf tali, operasi gabungan, pelabelan harmonis ganjil*

Abstract

A graph with the labeling properties of odd harmonic is called an odd harmonious graph. The purpose of this research was to get labeling properties of odd harmonic on the class of string graphs. The research used was a qualitative research method. The result of the research was that the definition and construction of a string graph, the union of a string graph, and the multiple string graph are obtained. Furthermore, it has been proved that a string graph, the union of a string graph, and the multiple string graph is an odd harmonious graph.

Keywords: *odd harmonious graph, odd harmonious labeling, multiple graph, string graph, union operation*

Diterima :

Direvisi :

Disetujui:

Copyright © 2019 : Barekeng: Jurnal ilmu matematika dan terapan, Jurusan Matematika Fakultas MIPA Universitas Pattimura, Ambon



1. INTRODUCTION

Graph labeling is one of the research topics on graph theory that has developed very rapidly in recent years. In 2019, Liang and Bai construct the definition of an odd harmonious graph as follows. For example, $G(p, q)$ is a graph with the set of vertices $V(G)$ and the set of lines $E(G)$ with order $p = |V(G)|$ and size $q = |E(G)|$. If the graph $G(p, q)$ fulfills the injective $f: V(G) \rightarrow \{0, 1, 2, 3, 4, 5, \dots, 2q - 1\}$ that induces $f^*: E(G) \rightarrow \{1, 3, 5, 7, 9, \dots, 2q - 1\}$ which is bijective with the definition of $f^*(x, y) = f(x) + f(y)$ so the graph $G(p, q)$ is an odd harmonious graph [1].

Gallian in 2019 collected all the research on graph labeling theory and its applications [2]. There are several classes of odd harmonious graphs that have been found, including: Srividya and Govindarajan in 2020 found a circular graph with parallel chord [3]. Ferbriana and Sugeng in 2020 found a squid graph [4], Jeyanthi and Philo in 2020 found a ladder graph [5]. Mumtaz et al in 2021 found a matting graph [6], Mumtaz and Silaban in 2021 found that the snake hair graph [7]. Pujiwati et al in 2021 found two star graphs [8]. Other relevant research results can be seen in [9], [10], [11], [12], [13], [14], [15], [16], [17], [18], [19], and [20].

In 2017, Firmansah and Yuwono found a snake net graph [21], and Firmansah, in 2020, found a multiple net snake graph which is an odd harmonious graph [22]. The two results in this paper are the basis for research to obtain new class constructions in the form of string graphs, combined string graphs, and multiple string graphs. This research aimed to find the labeling properties of odd harmonic in several classes of string graphs. The research method used was qualitative research. In this study, the construction and definition of the string graph $R(s)$ with $s \geq 1$, the combination of the string graph $R(s) \cup R(s)$ with $s \geq 1$, and the multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ will be given. Next, we will prove that the string graph $R(s)$ with $s \geq 1$, the combination of the string graph $R(s) \cup R(s)$ with $s \geq 1$, and the multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ fulfill the labeling properties of odd harmonious such that it is an odd harmonious graph. Furthermore, the results of this study also add new properties of odd harmonious graphs from previous studies.

2. RESEARCH METHOD

This research is a type of qualitative research that focuses on developing new properties of odd harmonious graphs. The research stages were as follows: Data collection stage, looking for relevant reference sources to obtain new graph class constructions in the form of graph class definitions and graph class images to obtain point notation and line notation from the graph class. The data analysis stage will be given the construction of a set of points, a set of lines, labeling points that are injective, and labeling lines that are bijective. Stages of theorem construction and proof, at this stage, the construction of the theorem and its proof will be given. Then, the theorem was proven using the direct proof method, namely by showing that the labeling function of the vertices that have been constructed fulfills the injective properties to induce a bijective arc labeling function.

3. RESULTS AND DISCUSSION

In this chapter, odd harmonious labeling will be given to several classes of string graphs, namely string graphs, combined string graphs and multiple string graphs.

3.1 Odd Harmonious Labeling on String Graphs

Definition 1. *string graph $R(s)$ with $s \geq 1$ is a graph with*

$$V(R(s)) = \{u_i | 0 \leq i \leq s + 1\} \cup \{v_i^j | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{w_i^j | 1 \leq i \leq s, j = 1, 2\} \text{ and}$$

$$E(R(s)) = \{u_i v_{(i+1)}^j | 0 \leq i \leq s, j = 1, 2\} \cup \{v_i^j u_i | 1 \leq i \leq s + 1, j = 1, 2\} \cup$$

$$\{v_i^j w_i^j | 1 \leq i \leq s, j = 1, 2\} \cup \{w_i^j v_{(i+1)}^j | 1 \leq i \leq s, j = 1, 2\}.$$

In Figure 1 given the construction of a string graph $R(s)$ with $s \geq 1$.




Figure 1. string graph $R(s)$

Next, we will prove that the string $R(s)$ with $s \geq 1$ is an odd harmonious graph expressed in Theorem 2.

Theorem 2. *The string graph $R(s)$ with $s \geq 1$ is odd harmonic*

Proof. Based on Definition 1 obtained $p = |V(R(s))| = 5s + 4$ and $q = |E(R(s))| = 8s + 4$. given $f: V(R(s)) \rightarrow \{0, 1, 2, 3, 4, 5, \dots, 16s + 7\}$

$$f(u_i) = 4i, 0 \leq i \leq s + 1 \tag{1}$$

$$f(v_i^j) = 4i + 2j - 5, j = 1, 2 \text{ and } 0 \leq i \leq s + 1 \tag{2}$$

$$f(w_i^j) = 16k - 12i - 4j + 18, j = 1, 2 \text{ and } 1 \leq i \leq s \tag{3}$$

Based on (1), (2), and (3) it is obtained that each point has a different table and $V(R(s)) \subseteq \{0, 1, 2, 3, 4, 5, \dots, 16s + 7\}$ so f is injective. Given $f^*: E(R(s)) \rightarrow \{1, 3, 5, 7, 9, \dots, 16s + 7\}$.

$$f^*(u_i v_{(i+1)}^j) = 2j + 8i - 1, j = 1, 2 \text{ and } 0 \leq i \leq s \tag{4}$$


$$f^*(v_i^j u_i) = 2j + 8i - 5, j = 1, 2 \text{ and } 1 \leq i \leq s + 1 \tag{5}$$

$$f^*(v_i^j w_i^j) = 16k - 2j - 8i + 13, j = 1, 2 \text{ and } 1 \leq i \leq s \tag{6}$$

$$f^*(w_i^j v_{(i+1)}^j) = 16k - 2j - 8i + 17, j = 1, 2 \text{ and } 1 \leq i \leq s \tag{7}$$

According to (4), (5), (6) and (7) It is obtained that each point has a different label and $E(R(s)) = \{1, 3, 5, 7, 9, \dots, 16s + 7\}$ so f^* is bijective. Therefore, the string graph $R(s)$ with $s \geq 1$ is an odd harmonious graph \square

Here is an example of a string graph: $R(6)$ in Figure 2 which is an odd harmonious graph.



Gambar 2. Graf tali $R(6)$

3.2 Odd Harmonious Labeling on Concatenated String Graphs

Definition 3. *Combined string graphs $R(s) \cup R(s)$ with $s \geq 1$ is a graph with $V(R(s) \cup R(s)) = \{u_i | 0 \leq i \leq s + 1\} \cup \{v_i^j | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{w_i^j | 1 \leq i \leq s, j = 1, 2\} \cup \{x_i | 0 \leq i \leq s + 1\} \cup \{y_i^j | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{z_i^j | 1 \leq i \leq s, j = 1, 2\}$ and $E(R(s) \cup R(s)) = \{u_i v_{(i+1)}^j | 0 \leq i \leq s, j = 1, 2\} \cup \{v_i^j u_i | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{v_i^j w_i^j | 1 \leq i \leq s, j = 1, 2\} \cup \{w_i^j v_{(i+1)}^j | 1 \leq i \leq s, j = 1, 2\} \cup \{x_i y_{(i+1)}^j | 0 \leq i \leq s, j = 1, 2\} \cup \{y_i^j x_i | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{y_i^j z_i^j | 1 \leq i \leq s, j = 1, 2\} \cup \{z_i^j y_{(i+1)}^j | 1 \leq i \leq s, j = 1, 2\}$.*

In Figure 3, the combined construction of a string graph is given $R(s) \cup R(s)$ with $s \geq 1$.




Figure 3. Combined string graphs $R(s) \cup R(s)$

Furthermore, it will be proved that the combined string $R(s) \cup R(s)$ is an odd harmonious graph which is stated in Theorem 4..

Theorem 4. Combined string graphs $R(s) \cup R(s)$ with $s \geq 1$ is odd harmonic

Proof. Based on Definition 3 obtained $p = |V(R(s) \cup R(s))| = 10s + 8$ and $q = |E(R(s) \cup R(s))| = 16s + 8$. Given $f: V(R(s) \cup R(s)) \rightarrow \{0, 1, 2, 3, 4, 5, \dots, 32s + 15\}$.

$$f(u_i) = 4i, 0 \leq i \leq s + 1 \quad (8)$$

$$f(v_i^j) = 4i + 2j - 5, j = 1, 2 \text{ and } 0 \leq i \leq s + 1 \quad (9)$$

$$f(w_i^j) = 16k - 12i - 4j + 18, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (10)$$

$$f(x_i) = 4i + 2, 0 \leq i \leq s + 1 \quad (11)$$

$$f(y_i^j) = 16k + 4i + 2j + 1, j = 1, 2 \text{ and } 0 \leq i \leq s + 1 \quad (12)$$

$$f(z_i^j) = 16k - 12i - 4j + 20, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (13)$$

According to (8), (9), (10), (11), (12), and (13) it is obtained that each point has a different label and $V(R(s) \cup R(s)) \subseteq \{0, 1, 2, 3, 4, 5, \dots, 32s + 15\}$ so f is injective. Defined line labeling function $f^*: E(R(s) \cup R(s)) \rightarrow \{1, 3, 5, 7, 9, \dots, 32s + 15\}$.

$$f^*(u_i v_{(i+1)}^j) = 8i + 2j - 1, j = 1, 2 \text{ and } 0 \leq i \leq s \quad (14)$$

$$f^*(v_i^j u_i) = 8i + 2j - 5, j = 1, 2 \text{ and } 1 \leq i \leq s + 1 \quad (15)$$

$$f^*(v_i^j w_i^j) = 16k - 8i - 2j + 13, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (16)$$

$$f^*(w_i^j v_{(i+1)}^j) = 16k - 8i - 2j + 17, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (17)$$

$$f^*(x_i y_{(i+1)}^j) = 16k + 8i + 2j + 7, j = 1, 2 \text{ and } 0 \leq i \leq s \quad (18)$$

$$f^*(y_i^j x_i) = 16k + 8i + 2j + 3, j = 1, 2 \text{ and } 1 \leq i \leq s + 1 \quad (19)$$

$$f^*(y_i^j z_i^j) = 32k - 8i - 2j + 21, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (20)$$

$$f^*(z_i^j y_{(i+1)}^j) = 32k - 8i - 2j + 25, j = 1, 2 \text{ and } 1 \leq i \leq s \quad (21)$$

Based on (14), (15), (16), (17), (18), (19), (20), and (21) it is obtained that each line has a different label and $E(R(s) \cup R(s)) = \{1, 3, 5, 7, 9, \dots, 32s + 15\}$ so f^* is bijective. Therefore, the combined graph of $R(s) \cup R(s)$ with $s \geq 1$ is an odd harmonious graph \square

The following is an example of a string graph combination $R(4) \cup R(4)$ in Figure 4 which is an odd harmonious graph.




Figure 4. Combined string graphs $R(4) \cup R(4)$

3.3 Odd Harmonious Labeling on a Multiple String Graph

Definition 5. Multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ is a graph with $V(R(s, t)) = \{u_i | 0 \leq i \leq s + 1\} \cup \{v_i^j | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{w_l^{i,j} | 1 \leq i \leq s, 1 \leq l \leq t, j = 1, 2\}$ and $E(R(s, t)) = \{u_i v_{(i+1)}^j | 0 \leq i \leq s, j = 1, 2\} \cup \{v_i^j u_i | 1 \leq i \leq s + 1, j = 1, 2\} \cup \{v_i^j w_l^{i,j} | 1 \leq i \leq s, 1 \leq l \leq t, j = 1, 2\} \cup \{w_l^{i,j} v_{(i+1)}^j | 1 \leq i \leq s, 1 \leq l \leq t, j = 1, 2\}$.

In Figure 5, the construction of a multiplied string graph is given $R(s, t)$ with $s \geq 1$ and $t \geq 1$




Figure 5. Graph of a multiple string $R(s, t)$

Next, it will be proved that the graph of a multiple $R(s, t)$ with $s \geq 1$ and $t \geq 1$ is an odd harmonious graph expressed in Theorem 6.

Theorem 6. The multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ is odd harmonious

Proof. Based on Definition 5, it is obtained $p = |V(R(s, t))| = 2st + 5s + 4$ and $q = |E(R(s))| = 4st + 8s + 4$. Given $f: V(R(s, t)) \rightarrow \{0, 1, 2, 3, 4, 5, \dots, 8st + 16s + 7\}$.

$$f(u_i) = 4i, 0 \leq i \leq s + 1 \quad (23)$$

$$f(v_i^j) = 4i + 2j - 5, 0 \leq i \leq s + 1, j = 1, 2 \quad (24)$$

$$f(w_l^{i,j}) = (8t + 16)s + 8l - (8t + 12)i - 4j + 10, 1 \leq i \leq s, 1 \leq l \leq t + 1, j = 1, 2 \quad (25)$$

Based on (23), (24), (25) it is obtained that each point has a different label and $V(R(s, t)) \subseteq \{0, 1, 2, 3, 4, 5, \dots, 8st + 16s + 7\}$ so f is injective. Given $f^*: E(R(s)) \rightarrow \{1, 3, 5, 7, 9, \dots, 8st + 16s + 7\}$.

$$f^*(u_i v_{(i+1)}^j) = 8i + 2j - 1, 0 \leq i \leq s, j = 1, 2 \quad (26)$$

$$f^*(v_i^j u_i) = 8i + 2j - 5, 1 \leq i \leq s + 1, j = 1, 2 \quad (27)$$

$$f^*(v_i^j w_l^{i,j}) = (8t + 16)s + 8l - (8t + 8)i - 2j + 5, 1 \leq i \leq s, 1 \leq l \leq t, j = 1, 2 \quad (28)$$

$$f^*(w_l^{i,j} v_{(i+1)}^j) = (8t + 16)s + 8l - (8t + 8)i - 2j + 9, 1 \leq i \leq s, 1 \leq l \leq t + 1, j = 1, 2 \quad (29)$$

Based on (26), (27), (28) and (29) it is obtained that each line has a different label and $E(R(s)) = \{1, 3, 5, 7, 9, \dots, 8st + 16s + 7\}$ so f^* is bijective. Therefore, multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ is an odd harmonious graph \square

The following is an example of a multiplied string graph $R(4, 3)$ in Figure 6 which is an odd harmonious graph.




Figure 6. The graph of a multiple string $R(4, 3)$

Based on the results in Theorem 2, Theorem 4 and Theorem 6, it has been proved that the string graph $R(s)$ with $s \geq 1$, the combination of the string graph $R(s) \cup R(s)$ with $s \geq 1$, and the multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ are odd harmonious graphs. The results and discussion show that this research is a development of previous studies [21] and [22], namely the addition of family graph classes from odd harmonious graphs.

4. CONCLUSION

Based on the results and discussion, it is obtained that the construction of the definition of a string graph $R(s)$ with $s \geq 1$, a combination of a string graph $R(s) \cup R(s)$ with $s \geq 1$, and a multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$. Furthermore, it has been proved that the string graph $R(s)$ with $s \geq 1$, the combination of the string graph $R(s) \cup R(s)$ with $s \geq 1$, and the multiple string graph $R(s, t)$ with $s \geq 1$ and $t \geq 1$ fulfils the labeling properties of odd harmonious such that it is an odd harmonious graph.

This research can be continued by looking for odd harmonious labeling on the combination of multiple string graphs $R(s, t) \cup R(s, t)$ with $s \geq 1$ and $t \geq 1$.

REFERENCES

- [1] Z.-H. Liang and Z.-L. Bai, "On the odd harmonious graphs with applications," *J. Appl. Math. Comput.*, vol. 29, no. 1–2, pp. 105–116, Jan. 2009.
- [2] J. A. Gallian, "A Dynamic Survey of Graph Labeling," *Electron. J. Comb.* 18., 2019.
- [3] V. Srividya and R. Govindarajan, "On odd harmonious labeling of even cycles with parallel chords and dragons with parallel chords," in *International Journal of Computer Aided Engineering and Technology*, 2020, vol. 13, no. 4.
- [4] F. Febriana and K. A. Sugeng, "Odd harmonious labeling on squid graph and double squid graph," in *Journal of Physics: Conference Series*, 2020, vol. 1538, no. 1.
- [5] P. Jeyanthi and S. Philo, "Odd harmonious labeling of step ladder graphs," *Util. Math.*, vol. 115, 2020.
- [6] K. Mumtaz, P. John, and D. R. Silaban, "The odd harmonious labeling of matting graph," in *Journal of Physics: Conference Series*, 2021, vol. 1722, no. 1.
- [7] K. Mumtaz and D. R. Silaban, "The odd harmonious labeling of n hair-kC4-snake graph," in *Journal of Physics: Conference Series*, 2021, vol. 1725, no. 1.
- [8] D. A. Pujiwati, I. Halikin, and K. Wijaya, "Odd harmonious labeling of two graphs containing star," in *AIP Conference Proceedings*, 2021, vol. 2326.
- [9] F. Firmansah, "Pelabelan Harmonis Ganjil pada Gabungan Graf Ular dan Graf Ular Berlipat," ["Odd Harmonious Labeling on Combined Snake Graphs and Multiple Snake Graphs,"] in *Konferensi Nasional Penelitian Matematika dan Pembelajarannya (KNPMP I)an*, 2016, pp. 809–818.
- [10] F. Firmansah and M. W. Syaifuddin, "Pelabelan Harmonis Ganjil pada Amalgamasi Graf Kincir Angin Belanda," ["Odd Harmonious Labeling on Dutch Windmill Graph Amalgamation,"] *Fibonacci J. Mat. dan Pendidik. Mat.*, 2018.
- [11] F. Firmansah and M. W. Syaifuddin, "Pelabelan Harmonis Ganjil pada Amalgamasi Graf Kincir Angin Double Quadrilateral," ["Odd Harmonious Labeling on Double Quadrilateral Windmill Graph Amalgamation,"] in *Seminar Nasional Pendidikan Matematika Ahmad Dahlan*, 2018, vol. 6.
- [12] F. Firmansah and Tasari, "Odd Harmonious Labeling on Edge Amalgamation from Double Quadrilateral Graphs," *Desimal J. Mat.*, vol. 3, no. 1, pp. 65–72, 2020.
- [13] P. Jeyanthi, S. Philo, and M. K. Siddiqui, "Odd harmonious labeling of super subdivision graphs," *Proyecciones*, vol. 38, no. 1, pp. 1–11, 2019.
- [14] J. Renuka and P. Balaganesan, "Odd harmonious labeling of some classes of cycle related graphs," *Indian J. Public Heal. Res. Dev.*, vol. 9, no. 9, pp. 403–408, 2018.
- [15] G. A. Saputri, K. A. Sugeng, and D. Froncek, "The odd harmonious labeling of dumbbell and generalized prism graphs," *AKCE Int. J. Graphs Comb.*, vol. 10, no. 2, pp. 221–228, 2013.
- [16] P. Jeyanthi, S. Philo, and K. A. Sugeng, "Odd harmonious labeling of some new families of graphs," *SUT J. Math.*, 2015.
- [17] P. Jeyanthi and S. Philo, "Odd harmonious labeling of some cycle related graphs," *Proyecciones*, vol. 35, no. 1, pp. 85–98, Mar. 2016.
- [18] F. Firmansah and M. R. Yuwono, "Odd Harmonious Labeling on Pleated of the Dutch Windmill Graphs," *Cauchy J. Mat. Murni dan Apl.*, vol. 4, no. 4, pp. 161–166, May 2017.
- [19] F. Firmansah, "Pelabelan Harmonis Ganjil pada Graf Bunga Double Quadrilateral," ["Odd Harmonious Labeling in Double Quadrilateral Interest Graphs,"] *J. Ilm. SAINS*, vol. 20, no. 1, 2020.
- [20] E. A. Pramesti and Purwanto, "Odd harmonious labeling of $S_n(m, r)$ graph," in *Journal of Physics: Conference Series*, 2021, vol. 1872, no. 1.
- [21] F. Firmansah and M. R. Yuwono, "Pelabelan Harmonis Ganjil pada Kelas Graf Baru Hasil Operasi Cartesian Product," ["Odd Harmonious Labeling on a New Graph Class Resulting from Cartesian Product Operations,"] *J. Mat. "MANTIK"*, vol. 3, no. 2, pp. 87–95, Oct. 2017.
- [22] F. Firmansah, "Pelabelan Harmonis Ganjil pada Graf Ular Jaring Berlipat," ["Odd Harmonious Labeling on Multiple-Net Snake Graphs,"] *Sainmatika J. Ilm. Mat. dan Ilmu Pengetah. Alam*, vol. 17, no. 1, p. 1, 2020.

ODD HARMONIOUS LABELING ON SOME STRING GRAPH CLASSES

ORIGINALITY REPORT

3%

SIMILARITY INDEX

PRIMARY SOURCES

1	ejournal.radenintan.ac.id Internet	32 words — 1%
2	ojs3.unpatti.ac.id Internet	26 words — 1%
3	seminar.uny.ac.id Internet	16 words — 1%

EXCLUDE QUOTES ON

EXCLUDE SOURCES OFF

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 15 WORDS